

Stable bundles on positive principal elliptic fibrations

Misha Verbitsky¹

verbit@maths.gla.ac.uk, verbit@mccme.ru

Abstract

Let $\pi \rightarrow X$ be a principal elliptic fibration over a Kähler base X . We assume that the Kähler form on X is lifted to an exact form on M (such fibrations are called **positive**). Examples of these are regular Vaisman manifolds (in particular, the regular Hopf manifolds) and Calabi-Eckmann manifolds. Assume that $\dim M > 2$. Using the Kobayashi-Hitchin correspondence, we prove that all stable bundles on M are flat on the fibers of the elliptic fibration. This is used to show that all stable vector bundles on M take form $L \otimes \pi^* B_0$, where B_0 is a stable bundle on X , and L a holomorphic line bundle. For X algebraic this implies that all holomorphic bundles on M are filtrable (that is, obtained by successive extensions of rank-1 sheaves). We also show that all positive-dimensional compact subvarieties of M are pullbacks of complex subvarieties on X .

Contents

1	Introduction	2
2	Principal elliptic fibrations	4
2.1	Positive elliptic fibrations: definition and examples	4
2.2	Preferred Hermitian metrics	5
3	Stable bundle on Hermitian manifolds	6
3.1	Gauduchon metrics and stability	6
3.2	Kobayashi-Hitchin correspondence	8
4	Hermitian-Einstein bundles on positive principal elliptic fibrations	8
4.1	Preferred metrics are Gauduchon	8
4.2	Primitive forms on positive principal elliptic fibrations	9
4.3	The Lübke-type positivity for Hermitian-Einstein bundles	10
4.4	Complex subvarieties in positive principal elliptic fibrations	11
4.5	Equivariance of stable bundles	11

¹Misha Verbitsky is an EPSRC advanced fellow supported by CRDF grant RM1-2354-MO02 and EPSRC grant GR/R77773/01

5 Line bundles on principal elliptic fibrations **12**

6 Structure theorem for stable bundles **14**

6.1 Equivariant \tilde{T} -action and the stable bundles 14

6.2 Filtrable coherent sheaves on positive elliptic fibrations 15

1 Introduction

Let M be a compact complex manifold, and $M \xrightarrow{\pi} X$ a smooth holomorphic fibration. Assume that an elliptic curve (considered as a complex Lie group) acts on M holomorphically. Assume, moreover, that this action is free and transitive on the fibers of π . Then $\pi : M \rightarrow X$ is called a **principal elliptic fibration**. The fibers of π are identified (non-canonically) with T .

This terminology is somewhat misleading, as in algebraic geometry one says “elliptic fibration” speaking of a fibration with a section. However, a principal bundle with a section is trivial, hence we use (following [BM2], [BM1]) the term “principal elliptic fibration” used to describe something which should be more properly called “a 1-dimensional torus principal bundle”.

In physics, the same object is usually called “a T^2 -bundle”. There is a great body of literature studying these manifolds in physical context, see e.g. [CCD], [GP] and the references therein. One usually considers a principal elliptic fibration over a 2-dimensional Calabi-Yau manifold, equipped with a Hermitian metric, such that its Bismut connection has holonomy $SU(3)$.¹ Such manifolds appear as a background for the heterotic string in the presence of fluxes.

Examples of principal elliptic fibrations are very common in complex geometry (take, for instance, the Hopf surface $\mathbb{C}^2 \setminus \{0\} / \langle q \rangle$, fibered over $\mathbb{C}P^1$ with a fiber $\mathbb{C}^* / \langle q \rangle$, where $q \in \mathbb{C}$, $|q| > 1$). For more examples see Subsection 2.1. The Hopf surface M is clearly homeomorphic to $S^1 \times S^3$. Since $H^2(M) = 0$, it is not Kähler. This is a general phenomenon. Unless π is trivialized after taking a finite cover, the total space M of a principal elliptic fibration is not Kähler.

In this paper we are interested in the so-called **positive principal elliptic fibrations** (Definition 2.1). These are the fibrations with the Kähler base X , such that some Kähler form ω_X on X is lifted to an exact form on

¹Bismut connection on M is a Hermitian connection in TM , with totally antisymmetric torsion; it is not difficult to show that such connection is unique on any Hermitian complex manifold.

M . Heuristically this means that the fibration $M \xrightarrow{\pi} X$ is positive in the sense of complex algebraic geometry (see Remark 2.4 for a more detailed explanation). From the positivity one obtains that M admits an exact positive 2-form $\pi^*\omega_X$ with all eigenvalues strictly positive except one.

In [Ve], [OV2] such form was used in Vaisman geometry to obtain Kodaira-type embedding theorem, several Kodaira-Nakano-type vanishing results and a study of compact subvarieties of a Vaisman manifold. We apply a similar reasoning to positive elliptic fibrations. Using the Kobayashi-Hitchin correspondence (Subsection 3.2), we obtain the following theorem.

Theorem 1.1: Let $M \xrightarrow{\pi} X$, $\dim_{\mathbb{C}} M = n$ be a positive principal elliptic fibration, (B, ∇) a Hermitian-Einstein bundle on M , and $\Theta \in \Lambda^{1,1}(M) \otimes \text{End}(B)$ its curvature. Assume that $n \geq 3$. Then $\Theta(v, \cdot) = 0$ for any vertical tangent vector $v \in T_{\pi}M$.

Proof: This is Theorem 4.3. ■

For a definition of stability and Hermitian-Einstein connections see Section 3. Theorem 1.1 implies the following corollary.

Proposition 1.2: Let T be an elliptic curve, and $M \xrightarrow{\pi} X$ a positive principal T -fibration, equipped with a preferred Hermitian metric. The universal covering \tilde{T} acts on M in a standard way (its action is factorized through T). Consider a stable bundle B on M . Then B is equipped with a natural holomorphic \tilde{T} -equivariant structure.

Proof: This is Proposition 4.6. ■

A similar argument proves

Proposition 1.3: Let $M \xrightarrow{\pi} X$ be a positive principal elliptic fibration, and $Z \subset M$ a closed positive-dimensional subvariety. Then $Z = \pi^*(Z_0)$ for some complex subvariety $Z_0 \subset X$.

Proof: This is Proposition 4.5. ■

Applying the \tilde{T} -equivariant structure arising on stable bundles, the following structure theorem is obtained

Theorem 1.4: Let $M \xrightarrow{\pi} X$ be a positive principal elliptic fibration

equipped with a preferred Hermitian metric, and B a stable holomorphic bundle on M . Then $B \cong L \otimes \pi^* B_0$, where L is a line bundle on M and B_0 a stable bundle on X .

Proof: This is Theorem 6.1. ■

We also show that all coherent sheaves on M are filtrable, that is, obtained as successive extensions of rank 1 sheaves (Theorem 6.5).

2 Principal elliptic fibrations

2.1 Positive elliptic fibrations: definition and examples

Throughout this paper, M is a compact complex manifold, X a Kähler manifold, T an elliptic curve, and $M \xrightarrow{\pi} X$ a principal T -fibration

Definition 2.1: A fibration is called **positive** if the pullback $\pi^* \omega_X$ is exact, for some Kähler form ω_X on X .

Principal toric fibrations, their invariants, topology and Dolbeault cohomology are thoroughly analyzed in the excellent paper [Hö]. For our purposes, the most important examples are the following.

Example 2.2: Regular Vaisman manifolds are principal elliptic fibrations constructed the following way (see e.g. [DO], [OV2]). Take a projective manifold X , and let L be an ample line bundle on X . Consider the space $\tilde{M} := \text{Tot}(L^* \setminus 0)$, which is a total space of the dual bundle L^* without the zero section. Then \tilde{M} is a principal \mathbb{C}^* -bundle over X . Fix $q \in \mathbb{C}$, $|q| > 1$, and let $M := \tilde{M} / \sim_q$, be the quotient of \tilde{M} under the equivalence $v \sim qv$, $v \in L^* \setminus 0$. Then M is a principal elliptic bundle, with a fiber $T = \mathbb{C}^* / \langle q \rangle$. The positivity of M is elementary (see e.g. [Ve] or [OV2]).

Vaisman manifolds were studied in great detail by I. Vaisman (see e.g. [Va1]), under the name of generalized Hopf manifolds.

Remark 2.3: A special case of the above example is a regular Hopf manifold $M \cong S^{2n-1} \times S^1$, obtained as a quotient of $\mathbb{C}^n \setminus 0$ under an equivalence generated by $v \rightarrow qv$. Clearly, M is fibered over $\mathbb{C}P^{n-1}$, with a fiber $T = \mathbb{C}^* / \langle q \rangle$. Regular Hopf manifold is obtained if one applies the construction of Example 2.2 to $X = \mathbb{C}P^{n-1}$, $L = \mathcal{O}(1)$.

Remark 2.4: Taking an arbitrary line bundle L on a Kähler manifold X and applying the same construction as above, we obtain a principal toric bundle as well. It will be positive in the sense of Definition 2.1 if and only if the bundle L is positive, as one can easily see from the exact sequence associated with the fibration in [Hö]

$$H^1(T) \xrightarrow{\delta} H^2(X) \xrightarrow{i} H^2(M). \quad (2.1)$$

The second arrow of (2.1) is a standard pullback map, and the transgression δ maps one of the generators of $H^1(T)$ to zero, the other to $c_1(L)$.

Example 2.5: The Calabi-Eckmann manifolds $M \cong S^{2n+1} \times S^{2m+1}$ are fibered in standard way over $\mathbb{C}P^n \times \mathbb{C}P^m$ (the spheres S^{2n+1} , S^{2m+1} are fibered over $\mathbb{C}P^n$, $\mathbb{C}P^m$ by the way of the Hopf fibration). The complex structure on M is constructed as follows. Fix a number $\tau \in \mathbb{C} \setminus \mathbb{R}$. Consider the action of an additive group \mathbb{C} on $\mathbb{C}^{n+1} \setminus \{0\} \times (\mathbb{C}^{m+1} \setminus \{0\})$,

$$t(v_1, v_2) \longrightarrow (e^t v_1, e^{t\tau} v_2).$$

This action is clearly holomorphic, and the quotient space M is naturally identified with $S^{2n+1} \times S^{2m+1}$,¹ moreover, M is equipped with a natural holomorphic projection $M \xrightarrow{\pi} \mathbb{C}P^n \times \mathbb{C}P^m$. The fibers of π are identified with $\mathbb{C}/\langle 1, \tau \rangle$. It is not difficult to see that M is a principal elliptic fibration. Since $H^2(M) = 0$, M is obviously positive.

2.2 Preferred Hermitian metrics

Let $M \xrightarrow{\pi} X$ be a principal elliptic fibration. As always, we assume that M is compact. By Blanchard's Theorem, unless π is trivialized on a finite covering, M is non-Kähler (see e.g. [Hö]). If M is positive, the pullback of a Kähler metric becomes exact, hence π cannot be trivialized. Therefore, a positive principal elliptic fibration is never Kähler.

We shall always assume that M is equipped with a special kind of Hermitian metric, defined as follows.

Recall that the smooth surjective map $\pi : (M, g) \longrightarrow (X, g_X)$ of Riemannian manifolds is called **Riemannian submersion** if its differential $d\pi$

¹To see this isomorphism, take a Hermitian metric on \mathbb{C}^{n+1} , \mathbb{C}^{m+1} . Then $|e^t v_1| = |e^{t\tau} v_2| = 1$ if and only if $\operatorname{Re}(t) = \log |v_1|$, $\operatorname{Re}(\tau t) = \log |v_2|$. Since $\tau \notin \mathbb{R}$, such t exists and is clearly unique.

induces an isometry $d\pi : T_\pi M^\perp \longrightarrow TX$, where $T_\pi M$ denotes the space of vertical tangent vectors, and $T_\pi M^\perp$ is orthogonal complement.

Definition 2.6: Let T be an elliptic curve, and $M \xrightarrow{\pi} X$ a positive principal T -fibration. Consider a Kähler metric g_X on X , such that the pullback $\pi^*\omega_X$ of the corresponding Kähler form is exact. A Hermitian metric g on M is called **preferred** if g is T -invariant, and the projection $\pi : (M, g) \longrightarrow (X, g_X)$ is a Riemannian submersion.

Clearly, preferred Hermitian metrics always exist.

Remark 2.7: Let M be a regular Vaisman manifold from Example 2.2, and $\tau \in TM$ the real vector field which is tangent to the map $v \longrightarrow \lambda v$, $\lambda \in \mathbb{R}$, $v \in L^*$. Then M admits a Hermitian metric g for which τ acts by isometries, and, moreover, $\nabla_{LC}\tau = 0$ for the Levi-Civita connection associated with this metric. In terminology used in Vaisman geometry, g is called the **locally conformally Kähler Gauduchon metric**, and τ **the Lee field** ([DO]). From the condition $\nabla_{LC}\tau = 0$ one immediately obtains that $\pi : (M, g) \longrightarrow (X, g_X)$ is a Riemannian submersion (see e.g. [OV1]). By construction, τ is a generator of the Lie algebra of T . Since τ acts by isometries, g is T -invariant. We obtain that the locally conformally Kähler Gauduchon metric on a regular Vaisman manifold is a preferred one.

3 Stable bundle on Hermitian manifolds

In this section, we recall the necessary definitions and explain the Kobayashi-Hitchin correspondence on non-Kähler manifolds. We follow [LT1] and [LT2].

3.1 Gauduchon metrics and stability

Definition 3.1: Let (M, g) be a Hermitian manifold, and $\omega \in \Lambda^{1,1}(M)$ its Hermitian form. We say that g is a Gauduchon metric if $\partial^*\bar{\partial}^*\omega = 0$, or, equivalently, $\partial\bar{\partial}(\omega^{\dim_{\mathbb{C}} M - 1}) = 0$.

Recall that the metrics g, g' on M are called **conformally equivalent** if $g = fg'$, $f \in C^\infty M$.

Theorem 3.2: ([Ga]) Let (M, g) be a compact Hermitian manifold. Then there exists a unique Gauduchon metric g' which is conformally equivalent

to g .

■

Definition 3.3: Let M be a compact complex manifold equipped with a Gauduchon metric, and $\omega \in \Lambda^{1,1}(M)$ the corresponding Hermitian form. Consider a torsion-free coherent sheaf F on M . Denote by $\det F$ its determinant bundle. Pick a Hermitian metric ν on $\det F$, and let Θ be the curvature of the associated Chern connection. We define the degree of F as follows:

$$\deg F := \int_M \Theta \wedge \omega^{\dim_{\mathbb{C}} M - 1}.$$

This notion is independent from the choice of the metric ν . Indeed, if $\nu' = e^{\psi} \nu$, $\psi \in C^{\infty}(M)$, then the associated curvature form is written as $\Theta' = \Theta + \partial\bar{\partial}\psi$, and

$$\int_M \partial\bar{\partial}\psi \wedge \omega^{\dim_{\mathbb{C}} M - 1} = 0$$

because ω is Gauduchon.

If F is a Hermitian vector bundle, Θ_F its curvature, and the metric ν is induced from F , then $\Theta = \text{Tr}_F \Theta_F$. In Kähler case this allows one to relate degree of a bundle with the first Chern class. However, in non-Kähler case, the degree is not a topological invariant — it depends fundamentally on the holomorphic geometry of F . Moreover, the degree is not discrete, as in the Kähler situation, but takes values in continuum.

Further on, we shall see that one can in some cases construct a holomorphic structure of any given degree $\lambda \in \mathbb{R}$ on a fixed C^{∞} -bundle. In our examples, such holomorphic structures are constructed on a topologically trivial line bundle over a positive principal elliptic fibration (Remark 5.4).

Definition 3.4: Let F be a non-zero torsion-free coherent sheaf on M . Then $\text{slope}(F)$ is defined as

$$\text{slope}(F) := \frac{\deg F}{\text{rk } F}.$$

The sheaf F is called

- stable** if for all subsheaves $F' \subset F$, we have $\text{slope}(F') < \text{slope}(F)$
- semistable** if for all subsheaves $F' \subset F$, we have $\text{slope}(F') \leq \text{slope}(F)$
- polystable** if F can be represented as a direct sum of stable coherent sheaves with the same slope.

Remark 3.5: This definition of stability is “good” as most standard properties of stable and semistable bundles hold in this situation as well. In particular, all line bundles are stable; all stable sheaves are simple; the Jordan-Hölder and Harder-Narasimhan filtrations are well defined and behave in the same way as they do in the usual Kähler situation ([LT1], [Br]).

However, not all bundles are **filtrable**, that is, are obtained as successive extensions by coherent sheaves of rank 1. There are non-filtrable vector bundles e.g. on a non-algebraic K3 surface (Example 6.4).

3.2 Kobayashi-Hitchin correspondence

The statement of Kobayashi-Hitchin correspondence is translated to the Hermitian situation verbatim, following Li and Yau ([LY]).

Definition 3.6: Let B be a holomorphic Hermitian vector bundle on a Hermitian manifold M , and $\Theta \in \Lambda^{1,1}(M) \otimes \text{End}(B)$ the curvature of its Chern connection ∇ . Consider the operator $\Lambda : \Lambda^{1,1}(M) \otimes \text{End}(B) \rightarrow \text{End}(B)$ which is a Hermitian adjoint to $b \rightarrow \omega \otimes b$, ω being the Hermitian form on M . The connection ∇ is called **Hermitian-Einstein** (or **Yang-Mills**) if $\Lambda\Theta = \text{const} \cdot \text{Id}_B$.

Theorem 3.7: (Kobayashi-Hitchin correspondence) Let B be a holomorphic vector bundle on a compact complex manifold equipped with a Gauduchon metric. Then B admits a Hermitian-Einstein connection ∇ if and only if B is polystable. Moreover, the Hermitian-Einstein connection is unique.

Proof: See [LY], [LT1], [LT2]. ■

Remark 3.8: The bundle B is stable if and only if the corresponding Hermitian-Einstein bundle (B, ∇) cannot be decomposed onto a direct sum of sub-bundles. In this case, the Hermitian-Einstein metric is defined by the connection uniquely up to a constant multiplier.

4 Hermitian-Einstein bundles on positive principal elliptic fibrations

4.1 Preferred metrics are Gauduchon

Proposition 4.1: Let $M \xrightarrow{\pi} X$ be a positive principal elliptic fibration,

and g a preferred Hermitian metric. Then g is Gauduchon.

Proof: Consider the orthogonal decomposition $TM = T_\pi M \oplus T_\pi M^\perp$ of the tangent bundle onto its vertical component and its orthogonal complement. We write the Hermitian form ω as $\omega = \omega_0 + \omega_T$, where ω_0 vanishes on $T_\pi M$ and ω_T vanishes on $T_\pi M^\perp$. Since π is a Riemannian submersion, $\omega_0 = \pi^* \omega_X$, where ω_X is a Kähler form on X . This implies

$$\partial^* \bar{\partial}^* \omega_0 = \pi^* \partial^* \bar{\partial}^* \omega_X = 0.$$

On the other hand, $\partial^* \bar{\partial}^* \omega_T$ can be computed by restricting ω_T to the fibers $\pi^{-1}(x)$ of π and computing $\partial^* \bar{\partial}^* (\omega_T|_{\pi^{-1}(x)})$. Since ω_T is T -invariant, $\partial^* \bar{\partial}^* (\omega_T|_{\pi^{-1}(x)})$ also vanishes, and we have

$$\partial^* \bar{\partial}^* \omega = \partial^* \bar{\partial}^* \omega_0 + \partial^* \bar{\partial}^* \omega_T = 0.$$

■

4.2 Primitive forms on positive principal elliptic fibrations

Let B_0 be a vector bundle with connection, and η a B_0 -valued form. Then η is called **closed** if $\nabla \eta = 0$, where

$$\nabla : \Lambda^i(M) \otimes B \longrightarrow \Lambda^{i+1}(M) \otimes B$$

is the connection operator extended to forms by the Leibniz rule. The curvature form of a connection is closed, by Bianchi identity.

Further on in this section, we shall need the following result.

Proposition 4.2: Let $M \xrightarrow{\pi} X$ be a positive principal elliptic fibration, $n = \dim M \geq 3$, equipped with a preferred Hermitian metric, B a Hermitian bundle with connection, and $\Theta \in \Lambda^{1,1}(M) \otimes \text{End}(B)$ a closed (1,1)-form. Assume that Θ is primitive, that is, $\Lambda \Theta = 0$. Then $\Theta(v, \cdot) = 0$ for any vertical tangent vector $v \in T_\pi M$.

Proof: Since the connection ∇ is Hermitian, it preserves the natural real structure in $\Lambda^{1,1}(M) \otimes \text{End}(B)$, $\eta \otimes b \longrightarrow \bar{\eta} \otimes b^\perp$, where by b^\perp one understands the Hermitian adjoint endomorphism. Therefore, we may assume that Θ is real, with respect to this real structure.

Let $\theta, \theta_1, \dots, \theta_{n-1}$ be an orthonormal basis in $\Lambda^{1,0}(M)$, with $\theta \in T_\pi M$, $\theta_i \in T_\pi M^\perp$. Consider a decomposition

$$\Theta = \sum_{i \neq j} (\theta_i \wedge \bar{\theta}_j + \bar{\theta}_i \wedge \theta_j) \otimes b_{ij} + \sum_i (\theta_i \wedge \bar{\theta}_i) \otimes a_i \quad (4.1)$$

$$+ \sum_i (\theta \wedge \bar{\theta}_i + \bar{\theta} \wedge \theta_i) \otimes b_i + \theta \wedge \bar{\theta} \otimes a, \quad (4.2)$$

with $b_{ij}, b_i, a_i, a \in \mathfrak{u}(B)$ being skew-Hermitian endomorphisms of B .

Consider now the form $\omega_0 := \pi^* \omega_X$. This form is exact, positive, and has $n - 1$ strictly positive eigenvalues. Using the basis described above, we can write

$$\omega = \sqrt{-1} \left(\theta \wedge \bar{\theta} + \sum_i \theta_i \wedge \bar{\theta}_i \right), \quad \omega_0 = \sqrt{-1} \left(\sum_i \theta_i \wedge \bar{\theta}_i \right)$$

where ω is the Hermitian form of M . This follows directly from ω being a preferred Hermitian form.

Let $\Xi := \text{Tr}(\Theta \wedge \Theta)$. This is a closed (2,2)-form on M . Then (4.1) implies

$$(\sqrt{-1})^n \Xi \wedge \omega_0^{n-2} = \text{Tr} \left(- \sum b_i^2 + a \left(\sum a_i \right) \right)$$

On the other hand, $\sum a_i + a = \Lambda \Theta = 0$, hence

$$(\sqrt{-1})^n \Xi \wedge \omega_0^{n-2} = \text{Tr} \left(- \sum b_i^2 - a^2 \right).$$

Since $\text{Tr}(-a^2)$ is a positive definite form on $\mathfrak{u}(B)$, the integral

$$\int_M (\sqrt{-1})^n \Xi \wedge \omega_0^{n-2} \quad (4.3)$$

is non-negative, and positive unless b_i and a both vanish everywhere. Using $n > 2$, we find that (4.3) vanishes, because ω_0 is exact and Ξ is closed. Therefore, b_i and a are identically zero, which is exactly the claim of Proposition 4.2

■

4.3 The Lübke-type positivity for Hermitian-Einstein bundles

Theorem 4.3: Let $M \xrightarrow{\pi} X$, $\dim_{\mathbb{C}} M = n$ be a positive principal elliptic fibration, equipped with a preferred Hermitian metric, (B, ∇) a Hermitian-Einstein bundle on M , and $\Theta \in \Lambda^{1,1}(M) \otimes \text{End}(B)$ its curvature. Assume that $n \geq 3$. Then $\Theta(v, \cdot) = 0$ for any vertical tangent vector $v \in T_\pi M$.

Proof: Theorem 4.3 is proven by the standard positivity argument, going back to M. Lübke ([Lü]). Similar argument is used e.g. to show that any Hermitian-Einstein bundle with vanishing Chern classes is flat.

Let $\Theta_0 := \Theta - \frac{1}{\text{rk } E} \text{Tr } \Theta$ be the traceless part of Θ . Then Θ_0 is primitive and closed, hence, by Proposition 4.2, $\Theta_0(v, \cdot) = 0$ for all $v \in T_\pi M$. To prove Theorem 4.3, it remains to show that the 2-form $\text{Tr } \Theta$ also vanishes on $v \in T_\pi M$. This is implied by the following trivial lemma.

Lemma 4.4: Let $M \xrightarrow{\pi} X$, $\dim_{\mathbb{C}} M \geq 3$ be a positive principal elliptic fibration, equipped with a preferred Hermitian metric, and $\eta \in \Lambda^{1,1}(M)$ a closed form satisfying $\Lambda\eta = \text{const}$. Then $\eta(v, \cdot) = 0$ for all $v \in T_\pi M$.

Proof: Clearly, $\Lambda\omega_0 = \text{const}$. Replacing η with $\eta - c\omega_0$, we may assume that $\Lambda\eta = 0$. Now η is primitive, and, by Proposition 4.2, $\eta(v, \cdot) = 0$ for all $v \in T_\pi M$. This finishes the proof of Theorem 4.3. ■

4.4 Complex subvarieties in positive principal elliptic fibrations

A similar argument proves the following result.

Proposition 4.5: Let $M \xrightarrow{\pi} X$ be a positive principal elliptic fibration, and $Z \subset M$ a closed positive-dimensional subvariety. Then $Z = \pi^*(Z_0)$ for some complex subvariety $Z_0 \subset X$.

Proof: The proof is taken from [Ve] verbatim, where the same result was proven for Vaisman manifolds.

It is well known that

$$\int_Z \omega_0^k \geq 0 \tag{4.4}$$

for all complex subvarieties $Z \subset M$, $\dim_{\mathbb{C}} Z = k$, and all positive forms ω_0 . Moreover, the integral (4.4) vanishes if and only if Z is tangent to the null-space foliation of ω_0 .

Since ω_0 is exact, the integral (4.4) vanishes. Therefore, Z is tangent to the null-space foliation of ω_0 . This implies Z is tangent to the fibers of π . ■

4.5 Equivariance of stable bundles

Theorem 4.3 has an immediate implication for the holomorphic geometry of

stable bundles.

Proposition 4.6: Let T be an elliptic curve, and $M \xrightarrow{\pi} X$, $\dim_{\mathbb{C}} M \geq 3$ a positive principal T -fibration, equipped with a preferred Hermitian metric. The universal covering \tilde{T} acts on M in a standard way (its action is factorized through T). Consider a stable bundle B on M . Then B is equipped with a natural holomorphic \tilde{T} -equivariant structure.

Proof: Let ∇ be a Hermitian-Einstein connection on B , and τ a holomorphic vector field tangent to the action of T . Consider the operator $\nabla_{\tau} : B \rightarrow B$. Since the curvature of B vanishes on τ , ∇_{τ} is holomorphic. For any $\lambda \in \mathbb{C}$ the operator $e^{\lambda \nabla_{\tau}}$ of parallel translation along τ defines a holomorphic \tilde{T} -equivariant structure on B . ■

Remark 4.7: The same argument implies that a stable bundle on a Vaisman manifold is always equivariant with respect to the action of the Lee field (see e.g. [DO], [OV2] for details).

5 Line bundles on principal elliptic fibrations

Let T be an elliptic curve and $M \xrightarrow{\pi} X$ a positive principal T -fibration, equipped with a preferred Hermitian metric. Any line bundle L on M is stable. From Proposition 4.6 we obtain a natural \tilde{T} -equivariant structure on L , and from Theorem 4.3 a connection ∇ which is flat on the fibers of π . Monodromy of this connection is related to the \tilde{T} -equivariant structure as follows. Let Γ be the kernel of the natural projection $\tilde{T} \rightarrow T$, $\Gamma = \pi_1 T \cong \mathbb{Z}^2$. Using the \tilde{T} -equivariant structure, we find that Γ acts on L by automorphisms. This action coincides with the monodromy of ∇ along the fibers of π .

The group of holomorphic automorphisms of L is identified with \mathbb{C}^* , therefore the monodromy map acts as a character $\chi : \Gamma \rightarrow \mathbb{C}^*$. Since Γ is a monodromy of a Hermitian connection, χ takes values in $U(1)$. We denote this character by $\chi_L : \Gamma \rightarrow U(1)$.

It turns out that any character can be realized by some line bundle. The following proposition is the main result of this section.

Proposition 5.1: Let T be an elliptic curve and $M \xrightarrow{\pi} X$ a positive principal T -fibration, $\dim M \geq 3$. Consider any character $\chi : \Gamma \rightarrow U(1)$. Then

there exists a holomorphic line bundle L on M such that the corresponding character $\chi_L : \Gamma \longrightarrow U(1)$ is equal to χ .

Proof: Consider the commutative diagram with exact rows coming from the exponential exact sequence

$$\begin{array}{ccccc} H^1(M, \mathcal{O}_M) & \longrightarrow & Pic_0(M) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \\ H^1(T, \mathcal{O}_T) & \longrightarrow & Pic_0(T) & \longrightarrow & 0 \end{array} \quad (5.1)$$

The characters $\chi : \Gamma \longrightarrow U(1)$ are in bijective correspondence with the bundles from $Pic_0(T)$, and the correspondence is provided by a unique flat connection on every $L \in Pic_0(T)$ (this follows e.g. from the Kobayashi-Hitchin correspondence on elliptic curve). Therefore, to prove Proposition 5.1 it is sufficient and necessary to show that the natural arrow $Pic_0(M) \longrightarrow Pic_0(T)$ is surjective. As one can see from (5.1), this is implied by surjectivity of the natural restriction map

$$H^1(M, \mathcal{O}_M) \longrightarrow H^1(T, \mathcal{O}_T). \quad (5.2)$$

Since $\dim H^1(T, \mathcal{O}_T) = 1$, it is actually sufficient to show that (5.2) is non-trivial. Looking again on (5.1), we find that to show non-triviality of (5.2) it is sufficient to prove that $Pic_0(M) \longrightarrow Pic_0(T)$ is non-trivial.

We reduced Proposition 5.1 to the following lemma

Lemma 5.2: In assumptions of Proposition 5.1, there exists a holomorphic line bundle L on M , $L \in Pic_0(M)$ such that L is non-trivial on fibers of $\pi : M \longrightarrow X$.

Remark 5.3: Restriction of L to different fibers of π is a flat bundle with the monodromy determined by the character χ_L . Therefore, the restriction $L|_{\pi^{-1}(x)}$ is independent from the choice of $x \in X$ (the fibers of π are naturally identified because the π is a principal fibration).

Proof of Lemma 5.2: Consider a trivial line bundle L_{triv} on M with trivial flat connection ∇_{triv} . Let θ be a 1-form which satisfies $d\theta = \omega_0$, where $\omega_0 = \pi^*\omega_X$ is the pullback of the Kähler form on X .

Consider the connection $\nabla_0 = \nabla_{triv} + \theta$ on L_{triv} . By definition, the curvature of ∇_0 is equal ω_0 . Therefore, $(\nabla_0^{0,1})^2 = 0$, and ∇_0 defines a

holomorphic structure on L_{triv} . Denote the corresponding holomorphic line bundle by L_0 . The degree of L_0 is easy to compute:

$$\deg L_0 = \int_M \omega \wedge \omega_0^{n-1} \quad (5.3)$$

and this number is clearly positive. Therefore, L_0 is a non-trivial holomorphic bundle. Lemma 5.2 is proven. We proved Proposition 5.1. ■

Remark 5.4: Let $q \in \mathbb{R}$ be a number. Consider the connection $\nabla' = \nabla_{triv} + q\theta$ on L_{triv} . Clearly, its curvature is equal $q\omega_0$. Therefore, ∇' defines a holomorphic structure on L_{triv} . Denote the corresponding holomorphic bundle by $L(q)$. From (5.3) we obtain that $\deg L(q) = q \deg L$. Therefore, the trivial C^∞ -bundle L_{triv} admits holomorphic structures of any given degree.

6 Structure theorem for stable bundles

6.1 Equivariant \tilde{T} -action and the stable bundles

The main result of this paper is the following theorem.

Theorem 6.1: Let $M \xrightarrow{\pi} X$, $\dim_{\mathbb{C}} M \geq 3$ be a positive principal elliptic fibration equipped with a preferred Hermitian metric, and B a stable holomorphic bundle on M . Then $B \cong L \otimes \pi^* B_0$, where L is a line bundle on M and B_0 a stable bundle on X .

Proof: As Proposition 4.6 implies, B is \tilde{T} -equivariant. The kernel Γ of the natural projection $\tilde{T} \rightarrow T$ acts on B by holomorphic automorphisms. Since B is stable, all its automorphisms are proportional to identity. Therefore, Γ acts on B by characters $\chi : \Gamma \rightarrow \mathbb{C}^*$. Since χ can be obtained via monodromy of the Hermitian-Einstein connection, χ takes values in $U(1)$. Let L be a line bundle on M which is \tilde{T} -equivariant and has the same monodromy action (such line bundle exists by Proposition 5.1). Then $B \otimes L^{-1}$ has trivial monodromy on the fibers of π . Clearly, $B \otimes L^{-1} = \pi^* B_0$ for some holomorphic bundle B_0 on X . Since L is by construction Hermitian-Einstein, the same is true for $B \otimes L^{-1}$ and for B_0 (the last assertion is true because π is a holomorphic Riemannian submersion). We proved Theorem 6.1. ■

Remark 6.2: In [BS] S. Bando and Y.-T. Siu developed Kobayashi-Hitchin correspondence for reflexive coherent sheaves. Using these results, it is easy

to extend Theorem 6.1 for reflexive coherent sheaves. We obtain that any stable reflexive coherent sheaf F on M is isomorphic to $L \otimes \pi^* F_0$, where F_0 is a stable reflexive coherent sheaf on X .

6.2 Filtrable coherent sheaves on positive elliptic fibrations

Definition 6.3: Let Z be a complex variety. A coherent sheaf F on Z is called **filtrable** if and only if the following equivalent conditions are satisfied.

- (i) F is obtained as a successive extension of coherent sheaves of rank 1
- (ii) F admits a filtration

$$0 = F_0 \subset F_1 \subset F_2 \dots \subset F_N = F,$$

$$\text{with } \text{rk}(F_i/F_{i-1}) = 1.$$

On a quasiprojective variety Z , every coherent sheaf is filtrable. This is not true if Z is not algebraic, as the following example shows.

Example 6.4: Let M be a K3 surface with $\text{Pic}(M) = 0$ (generic non-algebraic K3 surfaces have $\text{Pic}(M) = 0$). Since all line bundles on M are trivial, no stable vector bundle B , $\text{rk } B > 1$ can be filtrable. However, the tangent bundle TM is stable, because it is Hermitian-Einstein. Therefore, TM is not filtrable.

Theorem 6.5: Let $M \xrightarrow{\pi} X$ be a positive principal elliptic fibration, $\dim M \geq 3$. Assume that X is projective. Then all coherent sheaves on M are filtrable.

Proof: Let F be a coherent sheaf on M . Using the Harder-Narasimhan filtration ([Br]), we reduce Theorem 6.5 to the case when F is semistable. Using the Jordan-Hölder filtration, we reduce it to the case when F is a stable reflexive sheaf. By Theorem 6.1 (see Remark 6.2), $F = \pi^* F_0 \otimes L$, where F_0 is a stable reflexive sheaf on X . Since X is projective, F_0 is filtrable. Then F is also filtrable. We proved Theorem 6.5. ■

Acknowledgements: I am grateful to Ruxandra Moraru who explained to me the geometry of holomorphic bundles on elliptic fibrations and gave me reference I needed, and to Tony Pantev, who told me about the applications

and physical background of this geometry. Tony also found an error in a preliminary version of this paper. My gratitude also to F. Bogomolov and D. Kaledin, who gave me important insights on the geometry of Bott-Samelson manifolds.

References

- [BS] Bando, S., Siu, Y.-T, *Stable sheaves and Einstein-Hermitian metrics*, In: Geometry and Analysis on Complex Manifolds, Festschrift for Professor S. Kobayashi's 60th Birthday, ed. T. Mabuchi, J. Noguchi, T. Ochiai, World Scientific, 1994, pp. 39-50.
- [BM1] Vasile Brinzanescu, Ruxandra Moraru, *Stable bundles on non-Kähler elliptic surfaces*, math.AG/0306192, 15 pages.
- [BM2] Vasile Brinzanescu, Ruxandra Moraru, *Twisted Fourier-Mukai transforms and bundles on non-Kähler elliptic surfaces*, math.AG/0309031, 13 pages.
- [Br] L. Bruasse, *Harder-Narasimhan filtration on non-Kähler manifolds*, Int. Journal of Maths, 12(5):579-594, 2001.
- [CCD] G. L. Cardoso, G. Curio, G. Dall'Agata, D. Lüst, P. Manousselis and G. Zoupanos, *Non-Kähler string backgrounds and their five torsion classes*, 33 pages, hep-th/0211118.
- [DO] S. Dragomir, L. Ornea, *Locally conformal Kähler geometry*, Progress in Mathematics, 155. Birkhäuser, Boston, MA, 1998.
- [Ga] P. Gauduchon, *La 1-forme de torsion d'une variété hermitienne compacte*, Math. Ann., 267 (1984), 495–518.
- [GP] E. Goldstein and S. Prokushkin, *Geometric model for complex non-Kähler manifolds with $SU(3)$ structure*, 21 pages, hep-th/0212307.
- [Hö] Höfer, Thomas, *Remarks on torus principal bundles*, J. Math. Kyoto Univ. **33** (1993), no. 1, 227–259.
- [LY] Li, Jun, and Yau, S.-T., *Hermitian Yang-Mills connections on non-Kähler manifolds*, in “Mathematical aspects of string theory” (S.T. Yau ed.), World Scientific Publ., London, 1987, pp. 560-573.
- [Lü] M. Lübke, *Chernklassen von Hermite-Einstein Vektorbündeln*, Math. Ann. **260** (1982), 133-141.
- [LT1] Lübke, M., Teleman, A., *The Kobayashi-Hitchin correspondence*, World Scientific Publishing Co., Inc., River Edge, NJ, 1995. x+254 pp
- [LT2] Lübke, M., Teleman, A., *The universal Kobayashi-Hitchin correspondence on Hermitian manifolds*, math.DG/0402341, 90 pages.

- [M] Moraru, R., *Integrable systems associated to a Hopf surface*, *Canad. J. Math.* **55** (2003), no. 3, 609–635.
- [OV1] L. Ornea, M. Verbitsky, *Structure theorem for compact Vaisman manifolds*, math.DG/0305259, also in: *Math. Res. Lett.*, **10** (2003), no. 5-6, 799-805
- [OV2] L. Ornea, M. Verbitsky, *Immersion theorem for Vaisman manifolds*, math.AG/0306077, 28 pages
- [Va1] I. Vaisman, *Generalized Hopf manifolds*, *Geom. Dedicata* **13** (1982), no. 3, 231–255.
- [Ve] M. Verbitsky, *Vanishing theorems for locally conformal hyperkähler manifolds*, 2003, math.DG/0302219, 41 pages.

MISHA VERBITSKY

UNIVERSITY OF GLASGOW, DEPARTMENT OF MATHEMATICS,
15 UNIVERSITY GARDENS, GLASGOW G12 8QW, SCOTLAND.

INSTITUTE OF THEORETICAL AND EXPERIMENTAL PHYSICS
B. CHEREMUSHKINSKAYA, 25, MOSCOW, 117259, RUSSIA

verbit@maths.gla.ac.uk, verbit@mccme.ru